

PCT

WORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau




INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

<p>(51) International Patent Classification ⁶ : A61K 31/19, 31/22</p>	<p>A2</p>	<p>(11) International Publication Number: WO 95/34302 (43) International Publication Date: 21 December 1995 (21.12.95)</p>
<p>(21) International Application Number: PCT/US95/07570 (22) International Filing Date: 14 June 1995 (14.06.95) (30) Priority Data: 08/260,648 16 June 1994 (16.06.94) US (71) Applicant: ALLERGAN, INC. [US/US]; 2525 Dupont Drive, P.O. Box 19534, Irvine, CA 92713-9534 (US). (72) Inventors: ADORANTE, Joseph, S.; 1546 Thorn Tree Lane, Irvine, CA 92715 (US). WOLDEMUSSE, Elizabeth; 40 Gullwing Avenue, Laguna Niguel, CA 92677 (US). RUIZ, Guadalupe; 1068 Gentle Drive, Corona, CA 91720 (US). (74) Agents: BARAN, Robert, J. et al.; Allergan, Inc., 2525 Dupont Drive, P.O. Box 19534, Irvine, CA 92713-9534 (US).</p>		<p>(81) Designated States: AU, CA, JP, European patent (AT, BE, CH, DE, DK, ES, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE). Published <i>Without international search report and to be republished upon receipt of that report.</i></p>

(54) Title: METHOD FOR REDUCING INTRAOCULAR PRESSURE IN THE MAMMALIAN EYE BY ADMINISTRATION OF CALCIUM CHELATORS

(57) Abstract

Pharmaceutical compositions and a method are disclosed for treating glaucoma and/or ocular hypertension in the mammalian eye by administering to the mammalian eye the pharmaceutical composition of the invention which contains as the active ingredient one or more compounds having calcium chelating activity. Examples of calcium chelating agents utilized in the pharmaceutical composition and method of treatment are as in formula (2) and lower alkyl and alkoxyalkyl esters thereof.



FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AT	Austria	GB	United Kingdom	MR	Mauritania
AU	Australia	GE	Georgia	MW	Malawi
BB	Barbados	GN	Guinea	NE	Niger
BE	Belgium	GR	Greece	NL	Netherlands
BF	Burkina Faso	HU	Hungary	NO	Norway
BG	Bulgaria	IE	Ireland	NZ	New Zealand
BJ	Benin	IT	Italy	PL	Poland
BR	Brazil	JP	Japan	PT	Portugal
BY	Belarus	KE	Kenya	RO	Romania
CA	Canada	KG	Kyrgyzstan	RU	Russian Federation
CF	Central African Republic	KP	Democratic People's Republic of Korea	SD	Sudan
CG	Congo	KR	Republic of Korea	SE	Sweden
CH	Switzerland	KZ	Kazakhstan	SI	Slovenia
CI	Côte d'Ivoire	LI	Liechtenstein	SK	Slovakia
CM	Cameroon	LK	Sri Lanka	SN	Senegal
CN	China	LU	Luxembourg	TD	Chad
CS	Czechoslovakia	LV	Latvia	TG	Togo
CZ	Czech Republic	MC	Monaco	TJ	Tajikistan
DE	Germany	MD	Republic of Moldova	TT	Trinidad and Tobago
DK	Denmark	MG	Madagascar	UA	Ukraine
ES	Spain	ML	Mali	US	United States of America
FI	Finland	MN	Mongolia	UZ	Uzbekistan
FR	France			VN	Viet Nam
GA	Gabon				

METHOD FOR REDUCING INTRAOCULAR PRESSURE
IN THE MAMMALIAN EYE BY ADMINISTRATION
OF CALCIUM CHELATORS

5

BACKGROUND OF INVENTION

Field of the Invention

10 The present invention is directed to pharmaceutical compositions, and primarily to topically applied ophthalmic compositions comprising as the active ingredient one or more compounds having the ability to chelate calcium ions, e.g. intracellular calcium ions. The pharmaceutical compositions are useful for reducing intraocular pressure in animals of the mammalian species. In another
15 aspect, the present invention is directed to administering such formulations and compositions to animals of the mammalian species (including humans) for reducing intraocular pressure in the eye.

Brief Description of the Prior Art

20

Glaucoma is an optical neuropathy associated with elevated intraocular pressures which are too high for normal function of the eye, and results in irreversible loss of visual function. It is estimated in medical science that glaucoma afflicts approximately 2 per cent of the
25 population over the age of forty years, and is therefore a serious health problem. Ocular hypertension, i.e. the condition of elevated intraocular pressure, which has not yet caused irreversible damage, is believed to represent the earliest phase of glaucoma. Many therapeutic agents have been devised and discovered in the prior art for the treatment or
30 amelioration of glaucoma and of the condition of increased intraocular pressure which precedes glaucoma. Other compounds known to be useful in treating intraocular pressure are disclosed in the following patents.

35

United States Patent No. 3,467,756 describes anti-glaucoma and intraocular hypotensive compositions which contain in an ophthalmic vehicle 10, 11-dihydro-5-(3-methylaminopropyl)-5, 10-epoxy-11-hydroxy-5H-dibenzo [a,d]cycloheptene or related derivatives.

5 United States Patent No. 4,197,301 describes ophthalmic compositions which contain 1-(4-amino-6,7-dimethoxy-2-quinazolinyl)-4-(2-furanylcarbonyl)piperazine, also known under the name "prazosin".

10 United States Patent No. 4,565,821 describes a method of topically administering certain dopamine antagonists to reduce ocular hypertension and to treat glaucoma.

United States Patent No. 4,886,815 describes a method for treating retinal edema by administration of dopaminergic antagonists to a patient suffering from such conditions.

15 United States Patent No. 5,066,664 describes 2-hydroxy-2-alkylphenylamino)-oxazolines and thiazolines as anti-glaucoma and vasoconstrictive agents.

20 United States Patent No. 5,091,528 describe 6 or 7-(2-imino-2-imidazolidine)-1,4-benzoxazines as α adrenergic agents useful for treating glaucoma.


The foregoing and other anti-glaucoma and ocular hypotensive compounds and agents of the prior art do not provide such treatment or cure for glaucoma and ocular hypertension which is satisfactory in all respects. Therefore, the pharmacological and related arts and sciences
25 continue searching for additional and better anti-glaucoma and ocular hypotensive agents.

1,2-bis(2-aminophenoxy)ethane-N,N,N',N'-tetraacetic acid (BAPTA) is a specific Ca^{2+} chelator that has been used to clamp extracellular Ca^{2+} to desired levels. On the other hand, the
30 acetoxymethyl ester of BAPTA (BAPTA-AM) the uncharged esterified form of the parent compound is used to clamp intracellular Ca^{2+} (Ca^{2+}_i). BAPTA-AM penetrates biological cell membranes and is hydrolyzed by intracellular esterases yielding the original charged impermeable form of the compound once again capable of
35 buffering/clamping Ca^{2+} . (This is reported in Molecular Probes:


Handbook of Fluorescent Probes and Research Chemicals; Richard P. Haugland 1992-1994; Section 20: Calcium Indicators, Chelators and Ionophores, pages 119-128.)

5 SUMMARY OF THE INVENTION

Surprisingly it has been discovered in accordance with the present invention that calcium chelating agents are effective as anti-glaucoma agents and as agents for reducing intraocular pressure, when such agents are applied to the mammalian eye in a pharmaceutical composition, preferably in a topical ophthalmic composition. Accordingly, the present invention relates to a method of treating glaucoma, or ocular hypertension by topically administering to the mammalian eye an ophthalmic composition which contain an effective amount of a calcium chelating agent. A preferred example of calcium chelating agents suitable as the active ingredients of the ophthalmic compositions of the invention are:



and esters, e.g. the lower alkyl and alkoxyalkyl esters thereof. Such esters may be represented by the general formula



wherein R is lower alkyl, e.g. an alkyl radical having from 1 to 6 carbon

5

O

atoms, or $\text{---R}^1\text{---O---C---R}^2$ wherein R^1 is a lower alkylene radical, e.g. an alkylene radical having from 1 to 6 carbon atoms, and R^2 is R, as defined above. R, R^1 and R^2 may be interrupted with O or N radicals as in alkyloxy alkyl and alkylaminoalkyl moieties, e.g. R^1 may be $\text{---CH}_2\text{---N(CH}_3\text{)CH}_2\text{CH}_2\text{---}$ and $\text{---CH}_2\text{---O---CH}_2\text{---CH}_2\text{---}$. Preferably, R^1 and R^2 will comprise from 1 to 4 carbon atoms, e.g. 1 carbon atom.

10

While not wishing to be bound by theory it is believed that calcium chelating agents, e.g. 1,2-bis (2-aminophenoxy) ethane-N, N, N', N'-tetraacetic acid (BAPTA) or the acetoxymethyl ester of BAPTA (BAPTA-AM), are useful for treating hypertensive glaucoma, because intracellular (Ca^{2+}_i) is fundamental in the activation/control and modulation of epithelial fluid secretion. Thus, clamping/lowering the level of Ca^{2+}_i in the ciliary epithelium, the tissue responsible for aqueous humor formation and a determinant of intraocular pressure (IOP), will reduce inflow and therefore IOP.

20

The ophthalmic compositions of the invention contain the active ingredient in a concentration range of approximately 0.0001 to 0.1 per cent weight by volume. The composition itself includes, in addition to the active ingredient, such excipients which are per se well known in the art for preparing ophthalmic compositions, particularly ophthalmic solutions. In accordance with the method of the invention the

25

ophthalmic compositions, preferably ophthalmic solutions are applied topically to the mammalian eye approximately 1 or 2 times daily.

BRIEF DESCRIPTION OF THE DRAWINGS

5 Figure 1 is a graph showing the effect of topical administration of the drug BAPTA-AM on the regulatory volume decrease (RVD) of a suspension of cultured human non-pigmented ciliary epithelial cells.

10 Figure 2 is a graph showing the effect of intracameral administration of two dosages of the drug BAPTA-AM on the intraocular pressure (IOP) in the rabbit eye.

15 Figure 3 is a graph showing the effect of intracameral administration of the drug BAPTA-AM, as compared to saline and dimethyl sulfoxide (DMSO) on the intraocular pressure (IOP) in the rabbit eye.

DETAILED DESCRIPTION OF THE INVENTION

20 The compounds which are utilized in accordance with the method of the present invention, and in the pharmaceutical compositions of the present invention, are calcium chelating agents. In this regard the term calcium chelating agent is defined as those compounds which complex with calcium ions under physiological conditions, e.g. in an aqueous media at a pH of from 6.5 to 7.8. Specific and preferred examples of calcium chelating agents which are utilized in accordance with the present invention are provided below.

25 Pharmaceutically acceptable salts of the calcium chelating agents can also be used in accordance with the present invention. A pharmaceutically acceptable salt may be any salt which retains the activity of the parent compound and does not impart any deleterious or untoward effect on the subject to which it is administered and in the context in which it is administered.

30 Such a salt may be derived from any organic or inorganic acid or base. The salt may be a mono or polyvalent ion. Of particular interest where the acid function is concerned are the inorganic ions, e.g. sodium, potassium, etc. Organic amine salts may be made with amines, particularly ammonium salts such as mono-, di- and trialkyl amines or

ethanol amines. Salts may also be formed with caffeine, tromethamine and similar molecules. It is only important that the cation of any salt of a calcium chelating agent utilized in the compositions or methods of this invention be exchangeable with intracellular calcium ions.

5 For reducing intraocular pressure in a mammalian eye, and particularly for treatment of glaucoma in humans suffering from that condition, the active compounds (or mixtures or salts thereof) are administered in accordance with the present invention to the eye admixed with an ophthalmically acceptable carrier. Any suitable, e.g.,
10 conventional, ophthalmically acceptable carrier may be employed. A carrier is ophthalmically acceptable if it has substantially no long term or permanent detrimental effect on the eye to which it is administered. Examples of ophthalmically acceptable carriers include water (distilled or deionized water) saline and other aqueous media. In accordance with
15 the invention, the active compounds are preferably soluble in the carrier which is employed for their administration, so that the active compounds are administered to the eye in the form of a solution. Alternatively, a suspension of the active compound or compounds (or salts thereof) in a suitable carrier may also be employed.

20 In accordance with the invention the active compounds (or mixtures or salts thereof) are administered in an ophthalmically acceptable carrier in sufficient concentration so as to deliver an effective amount of the active compound or compounds to the eye. Preferably, the ophthalmic, therapeutic solutions contain one or more of the active
25 compounds in a concentration range of approximately 0.0001% to approximately 0.1% (weight by volume) and more preferably approximately 0.0005% to approximately 0.1% (weight by volume).

Any method of administering drugs directly to a mammalian eye may be employed to administer, in accordance with the present
30 invention, the active compound or compounds to the eye to be treated. By the term "administering directly" is meant to exclude those general systemic drug administration modes, e.g., injection directly into the patient's blood vessels, oral administration and the like, which result in the compound or compounds being systemically available. The primary
35 effect on the mammal resulting from the direct administering of the

active compound or compounds to the mammal's eye is preferably a reduction in intraocular pressure. More preferably, the active useful compound or compounds are applied topically to the eye or are injected directly into the eye. Particularly useful results are obtained when the

5 compound or compounds are applied topically to the eye in an ophthalmic solution (ocular drops).

Topical ophthalmic preparations, for example ocular drops, gels or creams, are preferred because of ease of application, ease of dose delivery, and fewer systemic side effects, such as cardiovascular

10 hypotentions. An exemplary topical ophthalmic formulation is shown below in Table I. The abbreviation q.s. means a quantity sufficient to effect the result or to make volume.

TABLE I

15	<u>Ingredient</u>	<u>Amount(% W/V)</u>
	Active Compound in accordance with the invention,	about 0.0001 to about 0.1
20	Preservative	0-0.10
	Vehicle	0-40
	Tonicity Adjustor	1-10
	Buffer	0.01-10
	pH Adjustor	q.s. pH 4.5-7.5
25	antioxidant	as needed
	Purified Water	as needed to make 100%

Various preservatives may be used in the ophthalmic preparation

30 described in Table I above. Preferred preservatives include, but are not limited to, benzalkonium chloride, chlorobutanol, thimerosal, phenylmercuric acetate, and phenylmercuric nitrate. Likewise, various preferred vehicles may be used in such ophthalmic preparation. These vehicles include, but are not limited to, polyvinyl alcohol, povidone,

35 hydroxypropyl methyl cellulose, poloxamers, carboxymethyl cellulose, hydroxyethyl cellulose, and purified water.


Tonicity adjustors may be added as needed or convenient. ~~They~~ include, but are not limited to, salts, particularly sodium chloride, potassium chloride, mannitol, and glycerin, or any other suitable ophthalmically acceptable tonicity adjustor.

5 Various buffers and means for adjusting pH may be used so long as the resulting preparation is ophthalmically acceptable. Accordingly, buffers include but are not limited to, acetate buffers, citrate buffers, phosphate buffers, and borate buffers. Acids or bases may be used to adjust the pH of these formulations as needed.

10 In a similar vein, ophthalmically acceptable antioxidants include, but are not limited to, sodium metabisulfite, sodium thiosulfate, acetylcysteine, butylated hydroxyanisole, and butylated hydroxytoluene.

 The ophthalmic solution (ocular drops) may be administered to the mammalian eye as often as necessary to maintain an acceptable level
15 of intraocular pressure in the eye. In other words, the ophthalmic solution (or other formulation) which contains the calcium chelating agent as the active ingredient, is administered to the mammalian eye as often as necessary to maintain the beneficial hypotensive effect of the active ingredient in the eye. Those skilled in the art will recognize that
20 the frequency of administration depends on the precise nature of the active ingredient and its concentration in the ophthalmic formulation. Within these guidelines it is contemplated that the ophthalmic formulation of the present invention will be administered to the mammalian eye approximately once or twice daily.

25 Specific examples of calcium chelating agents which are used as the active effective ingredients in the ophthalmic compositions of the present invention are described and shown below:



EXAMPLES

5

The present invention is demonstrated by *in vitro* and *in vivo* data. In Figure 1, 20 μ M BAPTA-AM were found to totally depress the regulatory volume decrease (RVD) that occurs following hypotonic swelling of cultured human non-pigmented ciliary epithelial (NPE) cells. In this example, NPE cells were loaded in an isosmotic (290 mOsm) solution containing 20 μ M BAPTA-AM for 30 minutes prior to suspension in a hypotonic (198 mOsm) solution. Control cells were subjected to the same hypotonic solution but without prior loading with BAPTA. Changes in cell volume were measured using a Coulter Counter interfaced to a Coulter Channelyzer. It is noted that, following osmotic swelling, control cells regulate towards their original isosmotic volume while BAPTA-loaded cells remain swollen. The above findings indicate that intracellular BAPTA via chelation of Ca^{2+} inhibits solute and osmotically obliged H_2O efflux. Because the Ca^{2+} -dependent ion flux pathways activated following osmotic cell swelling of NPE cells are involved in aqueous secretion, BAPTA will inhibit aqueous humor formation and, thus, lower IOP.

In the *in vivo* studies normotensive rabbits were injected intracamerally with 5 or 20 μ M BAPTA-AM. Figures 2 and 3 shows that 20 μ M BAPTA lowered IOP by 4 to 6 mm of Hg within 6 hours. Taken together the above *in vitro* and *in vivo* experiments demonstrate that chelation of Ca^{2+} in the ciliary epithelium will reduce IOP.

Several modifications of the present invention may become readily apparent to those skilled in the art in light of the present disclosure. For example, acetylcholine-like esters of BAPTA or other prodrugs of calcium chelating agents will provide target-specific activity for lowering IOP. In general since acetylcholine esterases reside primarily in the ciliary body and retina, the acetylcholine ester of BAPTA or another calcium chelating agent should only be hydrolyzed in the above tissue/cell types.


The advantages of using acetylcholine esters of BAPTA or another calcium chelating agent are as follows: First, Ca^{2+} will only be chelated/buffered in the ciliary epithelium and retina. Thus, other ocular cell types will be left unperturbed. Second, extracellular Ca^{2+} which is critical for maintaining tight junction integrity in fluid transporting epithelia will remain unaltered since the acetylcholine ester of BAPTA will not be appreciably hydrolyzed in the absence of esterases. Third, since the retina including retinal ganglion cells will also contain BAPTA in its ionized form, large increases in retinal ganglion cell Ca^{2+} will be prevented. Because increases in retinal and in particular optic nerve Ca^{2+} are believed to play a deleterious role in the pathophysiology of glaucoma and neural degeneration, intracellular BAPTA in the above cell types will afford an additional neuroprotective effect.

In view of the above, it is clear that the scope of the present invention should be interpreted solely on the basis of the following claims, as such claims are read in light of the disclosure.

30

WHAT IS CLAIMED IS:

1. A pharmaceutical composition useful for reducing
 5 intraocular pressure in the eye of a mammal, the composition
 comprising as its active ingredient one or more compounds having
 calcium chelating activity.
2. The pharmaceutical composition of Claim 1 wherein the
 10 composition is an ophthalmic solution, adapted for administration to
 the eye of a mammal in the form of eye droplets.
3. The pharmaceutical composition of Claim 1 wherein the
 compound having calcium chelating activity is present in the
 15 concentration range of 0.0001 to 0.1 per cent by volume.
4. The pharmaceutical composition of Claim 2 wherein the
 compound having calcium chelating activity is present in the
 concentration range of 0.0001 to 0.1 per cent weight by volume.
- 20 5. The pharmaceutical composition of Claim 1 where the
 compound having calcium chelating activity is selected from the group
 consisting of compounds represented by the formula :



25 and lower alkyl and alkoxyalkyl esters thereof.

6. The pharmaceutical composition of Claim 5 wherein the composition contains approximately 0.0001 to 0.1 per cent weight by volume of said compound having calcium chelating activity.

5


7. The pharmaceutical composition of Claim 6 where the composition is an ophthalmic solution, adapted for administration to the eye of a mammal in the form of eye droplets.

10

8. A pharmaceutical composition useful for providing neuroprotective activity in the eye of a mammal comprising as its active ingredient one or more compounds having calcium chelating activity.

15

9. The pharmaceutical composition of claim 8 wherein the compound having calcium chelating activity is selected from the group consisting of compounds represented by the formula:



and lower alkyl and alkoxyalkyl esters thereof.

20

10. The pharmaceutical composition of claim 9 wherein the composition contains approximately 0.0001 to 0.1 percent weight by volume of said compound having calcium chelating activity.

25


11. A method of treating animals of the mammalian species, including humans, for the purpose of reducing intraocular pressure in the eye of the mammal, the method of treatment comprising the step of

administering to the mammal a pharmaceutical composition which comprises as its active ingredient one or more compounds having calcium chelating activity.

5 12. The method of treatment of Claim 11 where the composition is an ophthalmic solution adapted for administration to the eye of a mammal in the form of eye droplets.

10 13. The method of treatment of Claim 12 where in the ophthalmic composition the concentration of the compound having calcium chelating activity is in the range of approximately 0.0001 to 0.1 per cent weight by volume.

15 14. A method of treating animals of the mammalian species, including humans, for the purpose of reducing intraocular pressure in the eye of the mammal, the method of treatment comprising the steps of administering to the mammal an ophthalmic composition which comprises as its active ingredient one or more compounds having calcium chelating activity compounds selected from the group
20 consisting of compounds represented by the formula:



and lower alkyl and alkoxyalkyl esters thereof.


25

15. The method of treatment of Claim 14 wherein the composition is an ophthalmic solution adapted for administration to the eye of a mammal in the form of eye droplets.

5 16. The method of treatment of Claim 14 wherein in the ophthalmic composition the concentration of the compound having calcium chelating activity is in the range of approximately 0.0001 to 0.1 per cent weight by volume.

10 17. A method for providing neuroprotective effect to the eye of a mammal which comprises the step of administering to the mammal a pharmaceutical composition which comprises as its active ingredient one or more compounds having calcium chelating activity.

15 18. The method of claim 17 where the compound having calcium chelating activity is selected from the group consisting of compounds represented by the formula:




20

and lower alkyl and alkoxyalkyl esters thereof.

19. The method of claim 18 wherein the composition contains approximately 0.0001 to 0.1 per cent weight by volume of said compound
25 having calcium chelating activity.

20. The pharmaceutical composition of claim 1 wherein the compound having calcium chelating activity is selected from the groups consisting of compounds represented by the formula:




5

wherein R is selected from the group consisting of alkyl radicals having
 from 1 to 6 carbon atoms and $\text{---R}^1\text{---O---C---R}^2$ wherein R¹ is an alkylene
 radical having from 1 to 6 carbon atoms and R² is R and wherein R, R¹
 and R² may be interrupted with O or N radicals.

21. The composition of claim 20 wherein R is selected from the
 group consisting of $\text{---CH}_2\text{---O---C---CH}_3$ and $\text{---CH}_2\text{---N(CH}_3\text{)CH}_2\text{CH}_2\text{---C---CH}_3$.

22. The composition of claim 21 wherein R is $\text{CH}_2\text{---O---C---CH}_3$.

23. The composition of claim 8 wherein the compound having calcium chelating activity is selected from the groups consisting of compounds represented by the formula:




wherein R is selected from the group consisting of alkyl radicals having

- 5 from 1 to 6 carbon atoms and $\text{---R}^1\text{---O---C---R}^2$ wherein R¹ is an alkylene radical having from 1 to 6 carbon atoms and R² is R and wherein R, R¹ and R² may be interrupted with O or N radicals.

- 10 24. The composition of claim 23 wherein R is selected from the group consisting of $\text{---CH}_2\text{---O---C---CH}_3$ and $\text{---CH}_2\text{---N(CH}_3\text{)CH}_2\text{CH}_2\text{---C---CH}_3$.

- 15 25. The composition of claim 24 wherein R is $\text{CH}_2\text{---O---C---CH}_3$.

- 20 26. The method of claim 11 wherein the compound having calcium chelating activity is selected from the groups consisting of compounds represented by the formula:




wherein R is selected from the group consisting of alkyl radicals having
O

- 5 from 1 to 6 carbon atoms and $\text{---R}^1\text{---O---C---R}^2$ wherein R^1 is an alkylene radical having from 1 to 6 carbon atoms and R^2 is R and wherein R, R^1 and R^2 may be interrupted with O or N radicals.

- 10 27. The method of claim 26 wherein R is selected from the
O O
group consisting of $\text{---CH}_2\text{---O---C---CH}_3$ and $\text{---CH}_2\text{---N(CH}_3\text{)CH}_2\text{CH}_2\text{---C---CH}_3$.

- 15 28. The composition of claim 26 wherein R is $\text{CH}_2\text{---O---C---CH}_3$.
O

29. The method of claim 17 wherein the compound having calcium chelating activity is selected from the groups consisting of compounds represented by the formula:



20

wherein R is selected from the group consisting of alkyl radicals having



from 1 to 6 carbon atoms and $-\text{R}^1-\text{O}-\text{C}-\text{R}^2$ wherein R^1 is an alkylene
5 radical having from 1 to 6 carbon atoms and R^2 is R and wherein R, R^1
and R^2 may be interrupted with O or N radicals.

30. The composition of claim 29 wherein R is selected from the



10 group consisting of $-\text{CH}_2-\text{O}-\text{C}-\text{CH}_3$ and $-\text{CH}_2-\text{N}(\text{CH}_3)\text{CH}_2\text{CH}_2-\text{C}-\text{CH}_3$.



31. The composition of claim 30 wherein R is $\text{CH}_2-\text{O}-\text{C}-\text{CH}_3$.

15

20

25

30

35




FIG. 1.




FIG. 2.




FIG. 3.